

## GENERAL INDEX TO VOLUME XXXI

New scientific names of plants and the final members of new combinations are printed in bold-face type; synonyms and page numbers having reference to figures and plates, in *italics*; and previously published and all other matter, in ordinary type.<sup>1</sup>

### A

- Acerates Floridana*, 368  
 Adaptation in yeasts, 219  
*Ageratum lineare*, 265  
 Alkali Ridge, Utah, prehistoric maize collection from, 345, 350, 352  
 Alleles, mating type, Instability of the, in *Saccharomyces*, 203  
*Amauroascus*, 186  
 Ammerman, Elizabeth, see Elizabeth Ammerman Baltzer  
 Amphiploidy, see Polyploidy  
 Anderson, Edgar. Cytological observations on *Tripsacum dactyloides*, 317; Homologies of the ear and tassel in Zea Mays, 325; Maiz reventador, 301; Notes on variation in *Tithonia tubaeformis*, 239; The sources of effective germ-plasm in hybrid maize, 355; Two collections of prehistoric corn tassels from southern Utah, 345; and Dorothy Schregardus. A method for recording and analyzing variations of internode pattern, 241  
 Araceae of Panama, 1  
 Arachniotus, the genus, 173; cytological study of, 174; history of, 173; systematic position of, 188  
*Arachniotus aureus*, 173, 174, 186; *candidus*, 173, 174, 186; *citrinus*, 174, 186; *ruber*, 173, 174, 186; *trachyspermus*, 174, 186; *trisporus*, 173, 174, 187, 190, 192, 194, 196, 198, development and systematic position of, The, 173  
 Archeological collections of corn tassels, 304, 335, 345, 350, 352  
 Asclepiadaceae from tropical America, Miscellaneous new, 235  
 Asclepiads, Notes on some North American, 363  
*Asclepias*, 363; *decumbens*, 368; *fascicularis*, 364, 365; *Floridana*, 368; *galioides*, 365; *incarnata*, 363; *Linaria*, 364; *mexicana*, 363; *revoluta*, 368; *Rolfii*, 368; *subverticillata*, 365; *tuberosa*, 366; ssp. *interior*, 368, 370, ssp. *tuberosa*, 368, 370, ssp. *Rolfii*, 368; *tuberosa*, 366, 368, var. *decumbens*, 367, var. *flexuosa*, 368; *verticillata*, 365, var. *subverticillata*, 365

*Aspergillaceae*, 188

*Axillaria*, 382

### B

- Baltzer, Elizabeth Ammerman. A monographic study of the genus *Palafoxia* and its immediate allies, 249  
*Bromeliaceae* of Panama, 73  
 Butterflyweed, 366

### C

- Camphor, effect of, on copulation in *Saccharomyces cerevisiae*, 203, 205, 206  
*Campydorium*, 382  
 Chiasma frequencies in *Tripsacum*, 320  
 Chromosome numbers in: *Coelorachis*, 321; *Tripsacum dactyloides*, 319; *Manisuris*, 321; *Polygonatum*, 377, 403; *Rottboellia*, 321; *Zea*, 319  
 Clines, 369  
*Commelinaceae* of Panama, 138  
*Convallaria*, 382; *angustifolia*, 393; *biflora*, 393; *canaliculata*, 393; *commutata*, 401; *gigantea*, 402; *hirta*, 412; *parviflora*, 393; *Polygonatum*, 382; *pubescens*, 383  
 Copulation: in *Arachniotus trisporus*, 180, 182; in *Saccharomyces cerevisiae*, 203, effect of camphor on, 203, 205, 206  
 Corn tassels, prehistoric, from southern Utah, Two collections of, 345  
 Corn, see Maize  
 Crozier formation in *Arachniotus*, 173, 188  
*Ctenomyces*, 186; *serratus*, 188  
*Cynanchum cubense*, 236; *Haughtii*, 235; *Marsdenioides*, 235; *peraffine*, 236  
 Cytological observations: on *Arachniotus trisporus*, 174; on *Polygonatum*, 377; on *Tripsacum dactyloides*, 317

### D

- Dermatophytes*, 188  
 Diakinesis, *Tripsacum dactyloides* mother cells at, 320  
 Diploid yeast cultures: adaptation in, as contrasted with haploid, 231; criteria for distinguishing, 209

### E

- Ear and tassel in Zea Mays, Homologies of the, 325

<sup>1</sup> For Woodson & Schery's "Flora of Panama" only the plant families and new entities are included in the ANNALS Index, since a complete Index will be appended at the end of each volume of the "Flora."

Enzymatic adaptation in yeasts, 219  
 Eriocaulaceae of Panama, 65  
 Euasclepias, 363  
 Euchlaena, 325; mexicana, 307  
 Evallaria, 382

## F

Favotrichum ochraceum, 188  
 Fernald's view of Asclepias tuberosa problem, 366  
 Fertilization in Arachniotus trisporus, 181  
 Flora of Panama, Part II, Fasc. 3, 1  
 Florestina, 254; callosa, 255  
 Frequency distributions for two prehistoric collections of corn, 348, 349

## G

Gaillardia lanceolata, 263  
 Galactose fermentation in yeasts, 219  
 Germ-plasm in hybrid maize, The sources of effective, 355  
 Gymnoascaceae, 173, 188; crozier formation in, 173  
 Gymnoascus, 186; aureus, 173, 186; candidus, 173, 186; Reessii, 188; ruber, 173, 186; sudans, 173, 174, 177, 187

## H

Haploid yeast cultures, criteria for distinguishing, 209  
 Heiser, Charles B., Jr. Monograph of Psilostrophe, 279  
 Hershey's story of Lancaster Surecopper corn, 358  
 Homologies of the ear and tassel in Zea Mays, 325  
 Hotson's strain of Arachniotus trisporus, 173, 174, 188, 200  
 Hybrid maize, The sources of effective germ-plasm in, 355  
 Hymenopappus integrifolius, 263

## I

Inflorescence in Zea Mays; main features of male, 327, 346, 346, female, 325  
 Instability of the mating type alleles in Saccharomyces, 302, under selection pressure, 207  
 Internode pattern: A method for recording and analyzing variations of, 241; of Salvia, 243, 245; of Tradescantia, 243, 244; of Tripsacum dactyloides, 245, 246

## J

Jalisco: maize grown in, 301; ears of maize reventador from, 312, 314; Tithonia

tubaeformis in, 239  
 Kelly, Dr. Isabel, 301  
 Kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of yeast, A comparison of the, 219

## L

Lancaster Surecopper corn, history of, 358  
 Lemnaceae of Panama, 60  
 Liliaceous genus Polygonatum in North America, The, 373  
 Lindegren, Gertrude, Carl C. Lindegren and: Instability of the mating type alleles in Saccharomyces, 203  
 Lindegren, Carl C.: and Gertrude Lindegren. Instability of the mating type alleles in Saccharomyces, 203; S. Spiegelman and. A comparison of the kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of yeasts, 219  
 Lomaxeta, 262; verrucosa, 263

## M

Maiz reventador, 301, 305, 312, 314; distribution of, 306, 309; uses of, 311; varieties mixed with, 314  
 Maiz, see Maize  
 Maize: Basket Maker, 349; Bearpaw Dent, 335, 342; breeding, 333, 337, 355; ear-tassel correlation in, 326, 329, 342; genetic background of, 330; germ-plasm, Sources of effective, in hybrid, 355; history of, in the Southwest, 349; Homologies of ear and tassel in, 325; hybrid, 307, 355; inbred lines of, 326; Inbred "Os 420", 335, 342; Krug, 356; Lancaster Surecopper, 342, 355; maize chapote, 302, 312; Maiz reventador, 301; maize rosquera, 301; open-pollinated, 355, Yellow Dent, 335, 342, 355; Mexican, 301, 329, 350, 352; Pueblo, 342; plant color, 303; prehistoric corn tassels from southern Utah, Two collections of, 345, 350, 352; Reid, 356; tassel, morphology of, 327, 327, score sheet showing technical details of, 347; Yellow Dent, 335, 342, 357  
 Manisuris, chromosome numbers of, 321  
 Matelea serpens, 236  
 Mayacaceae of Panama, 62  
 Media: for growing Arachniotus trisporus, 184; for growing Saccharomyces, 222  
 Megatrachophyton equinum, 188; ferrugineum, 188

- Method for recording and analyzing variations of internode pattern, A, 241  
 Mexican corn, 301, 329; introduction in American Southwest, 350  
 Mexico: maize in, 301; distribution of maize reventador in, 309, map, showing, 306  
 Microsporon, 188; Audouini, 188; lanosum, 188  
 Milkweeds, 363  
 Moldenke, Harold N. Eriocaulaceae of Panama, 65  
 Monograph of Psilostrophe, 279  
 Monographic study of the genus Palafoxia and its immediate allies, A, 249  
 Myxotrichum uncinatum, 188

## N

- Nannizzi's strain of Arachnitus trisporus, 174, 187, 188, 200  
 Natural selection, mathematical theory of, 223; applied to yeasts, 226  
 Notes on some North American Asclepiads, 363  
 Notes on variation in Tithonia tubaeformis, 239

## O

- Ownbey, Ruth Peck. The Liliaceous genus Polygonatum in North America, 373  
 Othake, 249, 253; callosum, 255, distribution of, 270; canescens, 257, distribution of, 272; Hookerianum, 261, distribution of, 274; Hookerianum, 260; Lindenii, 258, distribution of, 274; macrolepis, 258; maximum, 261; Reverchonii, 259, distribution of, 274; robustum, 257; roseum, 256, distribution of, 272; var. robustum, 257, distribution of, 272; sphacelatum, 259, distribution of, 274; tenuifolium, 255, 262; texanum, 257, distribution of, 274, var. macrolepis, 258, distribution of, 274

## P

- Palafoxia, A monographic study of Palafoxia and its immediate allies, 249  
 Palafoxia, 249, 264; arenaria, 266; callosa, 255; fastigiata, 263; Feayi, 267, distribution of, 276; Hookeriana, 260, 261, var. subradiata, 260; integrifolia, 263; leucophylla, 266; Lindenii, 258; linearis, 263, 270, distribution of, 276, var. arenicola, 266, var. gigantea, 266, distribution of, 276, var. leucophylla, 266, distribution of, 276; linearis, 266; Texana, 257, 261  
 Palafoxia, 253, 262

- Paleolaria, 262, 264; carnea, 265; fastigiata, 263  
 Panama, Flora of, Part II, Fasc. 3, 1  
 Pathogenicity of Arachnitus trisporus, 185  
 Peabody Museum of Harvard University, prehistoric corn tassels in, 345  
 Periballanthus, 382  
 Pinole, maize reventador used in, 311  
 Pioneer Hi-Bred Corn Company, tests with inbred lines of maize by, 333  
 Pipitillo, 301  
 Polygonatum, The Liliaceous genus, in North America, 373  
 Polygonatum, 382; angustifolium, 393; biflorum, 393, distribution of, 395, var. hebetifolium, 399, var. melleum, 399, var. necopinum, 401; biflorum, 383, 400, var. commutatum, 401, var. giganteum, 402, var. hebetifolium, 393,  $\gamma$ . latifolium, 383, var. ovatum, 402, var. ovatum subvar. ellipsoidale, 402, f. ramosum, 402, var. virginicum, 402; boreale, 384, var. australe, 384, var. multiflorum, 384; canaliculatum, 393, var. americanum, 402, var. americanum subvar. ellipticum, 402,  $\beta$ . giganteum, 402, var. giganteum, 402, var. oblongifolium, 402, var. ovatum, 402, f. ramosum, 402; cobrense, 392; commutatum, 401, distribution of, 404; commutatum var. lineamentosum, 402, var. ovatum, 402, f. ramosum, 402, var. virginicum, 402; cuneatum, 383; ellipticum, 402; Farwellii, 384; giganteum, 402; hebetifolium, 393, 400; hirtum, 412; latifolium, 402, var. commutatum, 401; melleum, 399; multiflorum,  $\beta$ . americanum, 383, 393, 402,  $\gamma$ . biflora, 393,  $\delta$ . canaliculata, 393,  $\gamma$ . giganteum, 402,  $\epsilon$ . latifolia, 402,  $\delta$ . latifolium, 383,  $\beta$ . pubescens, 383; officinale, 382; ovatum, 402; parviflorum, 393; pubescens, 383, distribution of, 385; pubescens var. australe, 384, var. boreale, 384, var. boreale subvar. australe, 384, var. cuneatum, 383, f. cuneatum, 384, f. fultius, 384, var. multiflorum, 384; virginicum, 402  
 Polyploidy in Tripsacum dactyloides, 317; in Polygonatum, 377  
 Polypteris, 249, 262; brasiliensis, 263; callosa, 253; Hookeriana, 260, 261; integrifolia, 262, 270, distribution of, 276; Lindenii, 258; macrolepis, 258; maxima, 261; Reverchonii, 259; robustum, 257; rosea, 256; Texana, 257  
 Polypteris 254

- Pontederiaceae of Panama, 151  
 Popcorn from maize reventador, 301  
 Populations of yeasts, 226  
 Prehistoric corn tassels from southern Utah,  
   Two collections of, 345, 350, 352  
 Psilostrophe, Monograph of, 279  
 Psilostrophe, 285; Bakeri, 288, 300, distribution of, 283; *biennis*, 293; *cerifera*, 293, var. *biennis*, 293; Cooperi, 287, 300, distribution of, 284; *divaricata*, 289, 290; *gnaphalodes*, 294, 300, distribution of, 283; *grandiflora*, 289, 292; *Hartmanii*, 290; *lanata*, 292; *pumila*, 288; *sparsiflora*, 289, 300, distribution of, 283; *sparsiflora*, 293; *tagetina*, 290, 300, distribution of, 284, var. *grandiflora*, 292, 300, distribution of, 284, var. *lanata*, 292, distribution of, 284, var. *sparsiflora*, 289; *Tagetinae*, 290, *villosa*, 293, 300, distribution of, 283  
 Pueblo corn, 349

## Q

- Quantitative inheritance in maize, 326, 339

## R

- Rapateaceae of Panama, 71  
 Reproduction, sexual, in *Arachniotus trisporus*, 180; ascus formation, 183, 198; copulation branches, 180, 182, 196; crozier formation, 183, 198; fertilization, 181  
*Riddellia*, 285; *arachnoidea*, 294; Cooperi, 287; *gnaphalioides*, 294; *tagetina*, 290, var. *pumila*, 288, var. *sparsiflora*, 289  
 Rosenbaum, Elisabeth Heuser. The development and systematic position of *Arachniotus trisporus*, 173

## S

- Saccharomyces cerevisiae*: Instability of the mating type alleles in, 203; galactozymase activity and natural selection in, 231  
 Sahuaripa, Relacion of (San Miguel de), manuscript copy of, 310  
*Salomonina*, 382; *biflora*, 393; *cobrensis*, 392; *commutata*, 402  
*Salvia*, internode pattern for species of, 245, 245  
 Schery, Robert W., Robert E. Woodson, Jr. and. Flora of Panama, Part II, Fasc. 3, 1  
 Schregardus, Dorothy, Edgar Anderson and. A method for recording and analyzing

- variations of internode pattern, 241  
*Sigillum*, 382  
*Siphyalis*, 382  
 Smith, Lyman B. Bromeliaceae of Panama, 73  
 Sources of effective germ-plasm in hybrid maize, The, 355  
 Spiegelman, S., and Carl C. Lindegren. A comparison of the kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of yeast, 219  
 Standley, Paul C. Araceae of Panama, 1  
*Stevia*, 253; *callosa*, 255; *linearis*, 265; *lavendulaefolia*, 265; *sphaecolata*, 259  
 Subspecies, three, of *Asclepias tuberosa*, 366

## T

- Tassels, maize: correlation between node condensation and row number, 333, 333; diagram showing main features of, 327, 346; morphology of, 326, 327; of maize reventador, 308; prehistoric, from southern Utah, Two collections of, 345; profiles, correlation between ear shape and, 326, 329, 336, of four inbred strains, 332; score sheet of, 347; spikelet arrangement in, 328  
 Temperature, optimum, for growth of *Arachniotus trisporus*, 185  
 Teosinte, 307  
*Tithonia tubaeformis*, Notes on variation in, 239  
 Torula: transformation of *Saccharomyces* into, 203; *utilis*, 203  
 Torulopsis, 203  
 Tradescantia, internode pattern for species of, 243, 244  
*Trichophyton*, 188; *asteroides*, 188; *denticulatum*, 188; *felineum*, 188; *radiolatum*, 188  
 Trichophytoneae, 188  
*Tripsacum dactyloides*: chiasma frequencies in, 320; chromosome numbers of, 319; Cytological observations on, 317; distribution of, 318; var. *occidentale*, 317; internode pattern of, 245, 246; pollen mother cells of, at diakinesis, 320

## U

- Utah, southern, Two collections of prehistoric corn tassels from, 345

## V

- Valonis' strain of *Arachniotus trisporus*, 173, 187, 200

Variation in *Tithonia tubaeformis*, Notes on, 239

Variations of internode pattern, A method for recording and analyzing, 241

## W

Woodson, Robert E., Jr.: Miscellaneous new Asclepiadaceae from tropical America, 235; Notes on some North American Asclepiads, 363; Robert W. Schery, and collaborators. Flora of Panama, Part II, Fasc. 3, 1

## X

Xyridaceae of Panama, 63

## Y

Yeast, A comparison of the kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of, 219

Yeasts: effect of camphor on copulation of, 203, 205, 206, 212, 214, 216; haploid and diploid, criteria for distinguishing, 209, nuclear fusion in, 210; instability of mating type alleles in, 203

## Z

Zea, chromosomes of, 319; Mays, Homologies of the ear and tassel in, 325

Zea Mays, see Maize

Zygosaccharomyces, 203



- Pontederiaceae of Panama, 151  
 Popcorn from maize reventador, 301  
 Populations of yeasts, 226  
 Prehistoric corn tassels from southern Utah,  
   Two collections of, 345, 350, 352  
 Psilostrophe, Monograph of, 279  
 Psilostrophe, 285; Bakeri, 288, 300, distribution of, 283; biennis, 293; cerifera, 293, var. biennis, 293; Cooperi, 287, 300, distribution of, 284; divaricata, 289, 290; gnaphalodes, 294, 300, distribution of, 283; grandiflora, 289, 292; Hartmanii, 290; lanata, 292; pumila, 288; sparsiflora, 289, 300, distribution of, 283; sparsiflora, 293; tagetina, 290, 300, distribution of, 284, var. grandiflora, 292, 300, distribution of, 284, var. lanata, 292, distribution of, 284, var. sparsiflora, 289; Tagetinae, 290, villosa, 293, 300, distribution of, 283  
 Pueblo corn, 349
- Q
- Quantitative inheritance in maize, 326, 339
- R
- Rapateaceae of Panama, 71  
 Reproduction, sexual, in Arachniotus trisporus, 180; ascus formation, 183, 198; copulation branches, 180, 182, 196; crozier formation, 183, 198; fertilization, 181  
 Riddellia, 285; arachnoidea, 294; Cooperi, 287; gnaphalioides, 294; tagetina, 290, var. pumila, 288, var. sparsiflora, 289  
 Rosenbaum, Elisabeth Heuser. The development and systematic position of Arachniotus trisporus, 173
- S
- Saccharomyces cerevisiae: Instability of the mating type alleles in, 203; galactozymase activity and natural selection in, 231  
 Sahuaripa, Relacion of (San Miguel de), manuscript copy of, 310  
 Salomonina, 382; biflora, 393; cobrensis, 392; commutata, 402  
 Salvia, internode pattern for species of, 245, 245  
 Schery, Robert W., Robert E. Woodson, Jr. and. Flora of Panama, Part II, Fasc. 3, 1  
 Schregardus, Dorothy, Edgar Anderson and. A method for recording and analyzing variations of internode pattern, 241  
 Sigillum, 382  
 Siphyalis, 382  
 Smith, Lyman B. Bromeliaceae of Panama, 73  
 Sources of effective germ-plasm in hybrid maize, The, 355  
 Spiegelman, S., and Carl C. Lindegren. A comparison of the kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of yeast, 219  
 Standley, Paul C. Araceae of Panama, 1  
 Stevia, 253; callosa, 255; linearis, 265; lavendulaefolia, 265; sphacelata, 259  
 Subspecies, three, of Asclepias tuberosa, 366
- T
- Tassels, maize: correlation between node condensation and row number, 333, 333; diagram showing main features of, 327, 346; morphology of, 326, 327; of maize reventador, 308; prehistoric, from southern Utah, Two collections of, 345; profiles, correlation between ear shape and, 326, 329, 336, of four inbred strains, 332; score sheet of, 347; spikelet arrangement in, 328  
 Temperature, optimum, for growth of Arachniotus trisporus, 185  
 Teosinte, 307  
 Tithonia tubaeformis, Notes on variation in, 239  
 Torula: transformation of Saccharomyces into, 203; utilis, 203  
 Torulopsis, 203  
 Tradescantia, internode pattern for species of, 243, 244  
 Trichophyton, 188; asteroides, 188; denticulatum, 188; felineum, 188; radiolatum, 188  
 Trichophytoneae, 188  
 Tripsacum dactyloides: chiasma frequencies in, 320; chromosome numbers of, 319; Cytological observations on, 317; distribution of, 318; var. occidentale, 317; internode pattern of, 245, 246; pollen mother cells of, at diakinesis, 320
- U
- Utah, southern, Two collections of prehistoric corn tassels from, 345
- V
- Valionis' strain of Arachniotus trisporus, 173, 187, 200

Variation in *Tithonia tubaeformis*, Notes on, 239

Variations of internode pattern, A method for recording and analyzing, 241

## W

Woodson, Robert E., Jr.: Miscellaneous new Asclepiadaceae from tropical America, 235; Notes on some North American Asclepiads, 363; Robert W. Schery, and collaborators. Flora of Panama, Part II, Fasc. 3, 1

## X

Xyridaceae of Panama, 63

## Y

Yeast, A comparison of the kinetics of enzymatic adaptation in genetically homogeneous and heterogeneous populations of, 219

Yeasts: effect of camphor on copulation of, 203, 205, 206, 212, 214, 216; haploid and diploid, criteria for distinguishing, 209, nuclear fusion in, 210; instability of mating type alleles in, 203

## Z

Zea, chromosomes of, 319; Mays, Homologies of the ear and tassel in, 325

Zea Mays, see Maize

Zygosaccharomyces, 203